



A service of the National Library of Medicine
and the National Institutes of Health

My NCBI
[Sign In] [Register]

All Databases

PubMed

Nucleotide

Protein

Genome

Structure

OMIM

PMC

Journals

Book

Search PubMed

for

Go

Clear

Limits

Preview/Index

History

Clipboard

Details

Display AbstractPlus

Show 20

Sort by

Send to

All: 1

Review: 0



☐ 1: Virology. 1999 Aug 15;261(1):8-14.

Full Text Article

Links

Structural analysis of adeno-associated virus transduction circular intermediates.

Duan D, Yan Z, Yue Y, Engelhardt JF.

Department of Anatomy and Cell Biology, College of Medicine, The University of Iowa, Iowa City 52242, USA.

Recombinant adeno-associated virus (rAAV) has recently been demonstrated to form circular intermediates following transduction in muscle tissue and cell lines. Although restriction enzyme and Southern blot analysis has revealed a consistent monomer and multimer head-to-tail conformation, detailed structural sequence analysis has been lacking due to the high secondary structure of the ITR arrays. To gain further insight into potential mechanisms by which AAV circular genomes are formed from linear single-stranded viral DNA, we have performed chemical sequencing of ITR arrays within seven circular intermediates independently isolated from primary fibroblasts and HeLa cells. Results from these studies demonstrated several types of circular intermediates with mosaic ITR elements flanked by two D sequences. The most predominant form consisted of a structure similar to that of previously generated AAV double-D plasmids, with one complete ITR flanked by two D-region elements. However, intermediately deleted ITR arrays with more than one complete ITR were also seen. Based on this structural information, we have proposed a model for formation of AAV circular intermediates by recombination/ligation between ITR ends of panhandle single-stranded AAV genomes.

PMID: 10484751 [PubMed - indexed for MEDLINE]

Related Links

Inverted terminal repeat sequences are important for intermolecular recombination and circularization of adeno-associated virus genome. [J Virol. 2005]

Consequences of DNA-dependent protein kinase catalytic subunit deficiency on recombinant adeno-associated virus genome circularization and heterodimerization in muscle tissue. [J Virol. 2005]

Cellular recombination pathways and viral terminal repeat hairpin structures are sufficient for adeno-associated virus integration in vivo and in vitro. [J Virol. 1997]

Circular intermediates of recombinant adeno-associated virus have defined structural characteristics responsible for long-term episomal persistence in muscle tissue. [J Virol. 1998]

Rescue and replication of adeno-associated virus type 2 as well as vector DNA sequences from recombinant plasmids containing deletions in the viral inverted terminal repeats: selective encapsidation of viral genomes in progeny virions. [J Virol. 1996]

See all Related Articles...

Display AbstractPlus

Show 20

Sort by

Send to

Write to the Help Desk

NCBI | NLM | NIH

Department of Health & Human Services

Privacy Statement | Freedom of Information Act | Disclaimer

Nov 6 2006 15:24:20